

LA-UR-14-21920

Approved for public release; distribution is unlimited.

Title: Application of the time-dependent close-coupling method to ionization of multi-electron atoms

Author(s): Armstrong, Gregory
Colgan, James P.

Intended for: Interview presentation at Auburn University

Issued: 2014-03-24



Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.



Application of the time-dependent close-coupling method to ionization of multi-electron atoms

Gregory Armstrong, James Colgan

Theoretical Division,
Los Alamos National Laboratory

UNCLASSIFIED

Plan of talk

- Introduction
- Two-photon double ionization of the Li
- Electron-impact ionization of Na and Mg
- Summary
- Future plans

UNCLASSIFIED

Introduction

- Time-dependent close-coupling (TDCC) method.
- Full-dimensionality numerical integration of the two/three-electron time-dependent Schrödinger equation.
- Applicable to a range of collision processes (photon, electron or ion impact).
- Accurate over a wide range of impact energies.

UNCLASSIFIED

Two-photon double ionization of lithium

UNCLASSIFIED

Motivation

- Provides guidance and support for recent measurements made at FLASH (Schuricke/Dorn).

UNCLASSIFIED

Motivation

- Provides guidance and support for recent measurements made at FLASH (Schuricke/Dorn).
- Such comparison requires accurate resolution of angular degrees of freedom and electron momenta.

UNCLASSIFIED

Motivation

- Provides guidance and support for recent measurements made at FLASH (Schuricke/Dorn).
- Such comparison requires accurate resolution of angular degrees of freedom and electron momenta.
- Provide first angular-resolved calculations for two-photon double ionization of Li.


UNCLASSIFIED

Motivation

- Provides guidance and support for recent measurements made at FLASH (Schuricke/Dorn).
- Such comparison requires accurate resolution of angular degrees of freedom and electron momenta.
- Provide first angular-resolved calculations for two-photon double ionization of Li.
- Application of TDCC codes to treatment of more than two photon absorptions (above-threshold ionization).

UNCLASSIFIED


FLASH experiments



- COLTRIMS (Cold Target Recoil Ion Momentum Spectroscopy) - Separate collection of ions and electrons.

UNCLASSIFIED


FLASH experiments



- COLTRIMS (Cold Target Recoil Ion Momentum Spectroscopy) - Separate collection of ions and electrons.
- Measured time of flight and position of ions/electrons allow momenta to be determined.

UNCLASSIFIED


FLASH experiments



- Two-photon double ionization of Li at 50 eV (direct) and 59 eV (intermediate excitation of the $1s2s2p$ state).

UNCLASSIFIED

FLASH experiments



- Two-photon double ionization of Li at 50 eV (direct) and 59 eV (intermediate excitation of the $1s2s2p$ state).
- $\text{Li } (1s^22s) + n\hbar\omega \rightarrow \text{Li}^{2+}(1s) + 2e^-$.

UNCLASSIFIED

Theory

UNCLASSIFIED

Time-dependent close-coupling method

- Full-dimensionality solution of the two-electron TDSE:

$$i \frac{\partial}{\partial t} \Psi(\mathbf{r}_1, \mathbf{r}_2, t) = H \Psi(\mathbf{r}_1, \mathbf{r}_2, t).$$

UNCLASSIFIED

Time-dependent close-coupling method

- Full-dimensionality solution of the two-electron TDSE:

$$i \frac{\partial}{\partial t} \Psi(\mathbf{r}_1, \mathbf{r}_2, t) = H \Psi(\mathbf{r}_1, \mathbf{r}_2, t).$$

- Finite-difference/basis-set expansion for each spin state:

$$\Psi^S(\mathbf{r}_1, \mathbf{r}_2, t) = \sum_{l_1, l_2, L} \frac{P_{l_1 l_2}^{LS}(r_1, r_2, t)}{r_1 r_2} |l_1 l_2 L\rangle.$$

UNCLASSIFIED

Time-dependent close-coupling method

- Full-dimensionality solution of the two-electron TDSE:

$$i \frac{\partial}{\partial t} \Psi(\mathbf{r}_1, \mathbf{r}_2, t) = H \Psi(\mathbf{r}_1, \mathbf{r}_2, t).$$

- Finite-difference/basis-set expansion for each spin state:

$$\Psi^S(\mathbf{r}_1, \mathbf{r}_2, t) = \sum_{l_1, l_2, L} \frac{P_{l_1 l_2}^{LS}(r_1, r_2, t)}{r_1 r_2} |l_1 l_2 L\rangle.$$

- Two active electrons move in the field of a frozen $1s \text{ Li}^{2+}$ orbital.

UNCLASSIFIED

TDCC method

- Angular reduction of the two-electron time-dependent Schrödinger equation yields

$$\begin{aligned}
 i \frac{\partial}{\partial t} P_{l_1 l_2}^{LS}(r_1, r_2, t) &= T_{l_1 l_2} P_{l_1 l_2}^{LS}(r_1, r_2, t) \\
 &+ \sum_{l'_1, l'_2} V_{l_1 l_2, l'_1 l'_2}^L P_{l'_1 l'_2}^{LS}(r_1, r_2, t) \\
 &+ \sum_{l'_1, l'_2, L'} W_{l_1 l_2, l'_1 l'_2}^{L, L'} P_{l'_1 l'_2}^{L' S}(r_1, r_2, t).
 \end{aligned}$$

UNCLASSIFIED

TDCC method

- Angular reduction of the two-electron time-dependent Schrödinger equation yields

$$\begin{aligned}
 i \frac{\partial}{\partial t} P_{l_1 l_2}^{LS}(r_1, r_2, t) &= T_{l_1 l_2} P_{l_1 l_2}^{LS}(r_1, r_2, t) \\
 &+ \sum_{l'_1, l'_2} V_{l_1 l_2, l'_1 l'_2}^L P_{l'_1 l'_2}^{LS}(r_1, r_2, t) \\
 &+ \sum_{l'_1, l'_2, L'} W_{l_1 l_2, l'_1 l'_2}^{L, L'} P_{l'_1 l'_2}^{L' S}(r_1, r_2, t).
 \end{aligned}$$

- Initial state obtained through relaxation of the field-free time-dependent Schrödinger equation in imaginary time.

UNCLASSIFIED

Double ionization of lithium at 50 eV

UNCLASSIFIED

Experimental considerations

- FLASH laser pulse length ~ 10 fs.
- Pulses display chaotic behaviour.
- Laser intensity will vary strongly over time.
- Intensity range: $5 \times 10^{13} \leq I \leq 5 \times 10^{15}$ (W/cm²).

UNCLASSIFIED

Triple-differential cross sections

UNCLASSIFIED

Triple-differential cross sections

- Momentum-space wavefunction, $\mathcal{P}_{l_1 l_2}^{LS}(\mathbf{k}_1, \mathbf{k}_2)$, obtained via projection of position-space wavefunction onto product of Li^{2+} continuum orbitals at the end of the laser pulse.

UNCLASSIFIED

Triple-differential cross sections

- Momentum-space wavefunction, $\mathcal{P}_{l_1 l_2}^{LS}(\mathbf{k}_1, \mathbf{k}_2)$, obtained via projection of position-space wavefunction onto product of Li^{2+} continuum orbitals at the end of the laser pulse.
- Appropriate integration of momentum-space wavefunction over all momenta yields the TDCS:

$$\frac{d^3\sigma}{dE_1 d\Omega_1 d\Omega_2} = \frac{1}{k_1 k_2} \left(\frac{\omega}{I}\right)^N \frac{1}{T_{\text{eff}}} \int_0^\infty dk_1 \int_0^\infty dk_2 \delta \left[\alpha - \tan^{-1} \left(\frac{k_2}{k_1} \right) \right] \times \sum_S w_S \left| \sum_{l_1, l_2, L} (-i)^{l_1+l_2} e^{i(\sigma_{l_1}+\sigma_{l_2})} \mathcal{P}_{l_1 l_2}^{LS}(k_1, k_2) |l_1 l_2 L\rangle \right|^2.$$

UNCLASSIFIED

Triple-differential cross sections


- Energy of outgoing electron pair may be partitioned arbitrarily for a given total excess energy.
- Coplanar detection geometry ($\phi_1 = \phi_2 = 0^\circ$) used to reduce angular variables from four to two.
- Polar angle of one electron is fixed, the other varied (relative to the laser polarization axis).

UNCLASSIFIED

Triple-differential cross sections

$1s2s\ ^1S$ initial state, equal energy sharing

$$\omega = 50\text{ eV}, I = 5 \times 10^{14}\text{ Wcm}^{-2}$$




- Anti-parallel emission also dominant for $\theta_1 = 0^\circ, 30^\circ$ and $\theta_1 = 150^\circ$, with additional near-perpendicular emissions.

UNCLASSIFIED

Triple-differential cross sections

$1s2s\ ^1S$ initial state, equal energy sharing

$$\omega = 50\text{ eV}, I = 5 \times 10^{14}\text{ Wcm}^{-2}$$




- Near-perpendicular emission dominant for $\theta_1 = 60^\circ$ and $\theta_1 = 120^\circ$.

UNCLASSIFIED

Triple-differential cross sections

$1s2s\ ^1S$ initial state, equal energy sharing

$$\omega = 50\text{ eV}, I = 5 \times 10^{14}\text{ Wcm}^{-2}$$




- Emission at a mutual angle $\theta_{12} = 120^\circ$ dominates for $\theta_1 = 90^\circ$.

UNCLASSIFIED

Triple-differential cross sections

$1s2s\ ^3S$ initial state, equal energy sharing

$$\omega = 50\text{ eV}, I = 5 \times 10^{14}\text{ Wcm}^{-2}$$




- Anti-parallel emission avoided for all values of θ_1 .

UNCLASSIFIED

Triple-differential cross sections

$1s2s\ ^3S$ initial state, equal energy sharing

$$\omega = 50\text{ eV}, I = 5 \times 10^{14}\text{ Wcm}^{-2}$$




- Dominant emission at $\theta_{12} \simeq 120^\circ$ at all angles.

UNCLASSIFIED

Triple-differential cross sections

1s2s 3S initial state, equal energy sharing

$$\omega = 50 \text{ eV}, I = 5 \times 10^{14} \text{ Wcm}^{-2}$$



- Two such contributions for $\theta_1 = 0^\circ$ and $\theta_1 = 90^\circ$.

UNCLASSIFIED

Recoil-ion momentum distributions

UNCLASSIFIED

Recoil-ion momentum distributions

- Recoil-ion momentum vector given in terms of electron momenta:

$$\mathbf{p} = -(\mathbf{k}_1 + \mathbf{k}_2) .$$

UNCLASSIFIED

Recoil-ion momentum distributions

- Recoil-ion momentum vector given in terms of electron momenta:

$$\mathbf{p} = -(\mathbf{k}_1 + \mathbf{k}_2) .$$

- Momentum distribution obtained via integration of TDCS over appropriate set of angles and energies

$$\frac{d^3\sigma}{dp_x dp_y dp_z} = \int d\Omega'_1 \int d\Omega'_2 \int dE_1 \frac{d^3\sigma}{d\Omega_1 d\Omega_2 dE_1} \delta(\mathbf{p} + \mathbf{k}_1 + \mathbf{k}_2) .$$


- Integration over all p_y gives distribution over p_x and p_z .

UNCLASSIFIED

Recoil-ion momentum distributions

Singlet and triplet combined

$$I = 5 \times 10^{14} \text{ Wcm}^{-2}$$




- Features at larger momenta ($|p_x, p_z| > 2$) indicate three-photon double ionization.

UNCLASSIFIED

Recoil-ion momentum distributions

Singlet and triplet combined

$$I = 1 \times 10^{15} \text{ Wcm}^{-2}$$




- Increase in magnitude of peak features associated with three-photon double ionization.

UNCLASSIFIED

Recoil-ion momentum distributions

Singlet and triplet combined

$$I = 5 \times 10^{15} \text{ Wcm}^{-2}$$



- Increase in central features ($p_z \simeq 0$) associated with anti-parallel emission.

UNCLASSIFIED


Comparison with experiment

UNCLASSIFIED

Comparison with experiment

TDCC

$$I = 1 \times 10^{15} \text{ Wcm}^{-2}$$



Experiment
(Schuricke/Dorn)

Singlet and triplet contributions

UNCLASSIFIED


Two photon absorptions

UNCLASSIFIED

Singlet contribution

Two photon absorptions

$$I = 1 \times 10^{15} \text{ Wcm}^{-2}$$




- Prominent central features due to anti-parallel emission.

UNCLASSIFIED

Triplet contribution

Two photon absorptions

$$I = 1 \times 10^{15} \text{ Wcm}^{-2}$$



- Central minimum - anti-parallel emission largely avoided.

UNCLASSIFIED

Double ionization of lithium at 59 eV

UNCLASSIFIED

Double ionization of lithium at 59 eV

- A minimum of two photons required for double ionization from an initial ground state.

UNCLASSIFIED

Double ionization of lithium at 59 eV

- A minimum of two photons required for double ionization from an initial ground state.
- Double ionization proceeds via initial photoexcitation of the $1s2s2p\ ^2P_{M=0}$ state.

UNCLASSIFIED

Double ionization of lithium at 59 eV

- A minimum of two photons required for double ionization from an initial ground state.
- Double ionization proceeds via initial photoexcitation of the $1s2s2p\ ^2P_{M=0}$ state.
- A single 59 eV photon is sufficient to ionize the 2s and 2p electrons.

UNCLASSIFIED

Double ionization of lithium at 59 eV

- A minimum of two photons required for double ionization from an initial ground state.
- Double ionization proceeds via initial photoexcitation of the $1s2s2p\ ^2P_{M=0}$ state.
- A single 59 eV photon is sufficient to ionize the 2s and 2p electrons.
- TDCC calculations model photoionization of $2s2p\ ^{1,3}P_{M=0}$ two-electron states.

UNCLASSIFIED


Triple-differential cross sections

UNCLASSIFIED

Triple-differential cross sections

2s2p 1P initial state, equal energy sharings

$$\omega = 59 \text{ eV}, I = 5 \times 10^{14} \text{ Wcm}^{-2}$$




- Dominant anti-parallel emission for $\theta_1 = 0^\circ$ with additional near-perpendicular components.

UNCLASSIFIED

Triple-differential cross sections

2s2p 1P initial state, equal energy sharings

$$\omega = 59 \text{ eV}, I = 5 \times 10^{14} \text{ Wcm}^{-2}$$




- Anti-parallel emission no longer dominant for $\theta_1 = 30^\circ$ and $\theta_1 = 150^\circ$.

UNCLASSIFIED

Triple-differential cross sections

2s2p 1P initial state, equal energy sharings

$$\omega = 59 \text{ eV}, I = 5 \times 10^{14} \text{ Wcm}^{-2}$$




- Near-perpendicular emission dominant for $\theta_1 = 60^\circ$ and $\theta_1 = 120^\circ$.

UNCLASSIFIED

Triple-differential cross sections

2s2p 1P initial state, equal energy sharings

$$\omega = 59 \text{ eV}, I = 5 \times 10^{14} \text{ Wcm}^{-2}$$




- Two peaks for $\theta_1 = 90^\circ$ at $\theta_2 = \theta_1 \pm 120^\circ$ with additional near-anti-parallel components.

UNCLASSIFIED

Triple-differential cross sections

2s2p 3P initial state, equal energy sharings

$$\omega = 59 \text{ eV}, I = 5 \times 10^{14} \text{ Wcm}^{-2}$$




- Anti-parallel emission avoided for all values of θ_1 .

UNCLASSIFIED

Triple-differential cross sections

2s2p 3P initial state, equal energy sharings

$$\omega = 59 \text{ eV}, I = 5 \times 10^{14} \text{ Wcm}^{-2}$$




- Dominant emission at $\theta_{12} \simeq 130^\circ$ at all values of θ_1 .

UNCLASSIFIED

Triple-differential cross sections

2s2p 3P initial state, equal energy sharings

$$\omega = 59 \text{ eV}, I = 5 \times 10^{14} \text{ Wcm}^{-2}$$



- Two such contributions for $\theta_1 = 0^\circ$ and $\theta_1 = 90^\circ$.


UNCLASSIFIED

Recoil-ion momentum distributions


UNCLASSIFIED

Comparison with experiment

(a) Experimental distribution



(b) Theoretical distribution




- Reasonable agreement in lobe structure positions and relative magnitudes.


UNCLASSIFIED

Singlet and triplet contributions

(c) Singlet contribution to (b)



(d) Triplet contribution to (b)




- Peaks at $p_z > 1$ in the singlet contribution indicate preference for emission into a common hemisphere.


UNCLASSIFIED

Singlet and triplet contributions

(c) Singlet contribution to (b)



(d) Triplet contribution to (b)




- Singlet contribution again shows central features due to anti-parallel emission.


UNCLASSIFIED

Singlet and triplet contributions

(c) Singlet contribution to (b)



(d) Triplet contribution to (b)



- No such features in the triplet contribution since anti-parallel emission is largely avoided.

UNCLASSIFIED

Summary

- First set of calculations for two- and three-photon double ionization of Li.

UNCLASSIFIED

Summary

- First set of calculations for two- and three-photon double ionization of Li.
- Each spin state shows highly characteristic emission configurations.

UNCLASSIFIED

Summary

- First set of calculations for two- and three-photon double ionization of Li.
- Each spin state shows highly characteristic emission configurations.
- Intermediate photoexcitation has a considerable effect on emission configurations.

UNCLASSIFIED

Summary

- First set of calculations for two- and three-photon double ionization of Li.
- Each spin state shows highly characteristic emission configurations.
- Intermediate photoexcitation has a considerable effect on emission configurations.
- Recoil-ion momentum distributions are in good agreement with FLASH measurements.

UNCLASSIFIED

Electron-impact ionization of Na and Mg

UNCLASSIFIED

Theory

UNCLASSIFIED

TDCC method

- Angular reduction of the two-electron time-dependent Schrödinger equation yields

$$i \frac{\partial}{\partial t} P_{l_1 l_2}^{LS}(r_1, r_2, t) = T_{l_1 l_2} P_{l_1 l_2}^{LS}(r_1, r_2, t) + \sum_{l'_1, l'_2} V_{l_1 l_2, l'_1 l'_2}^L P_{l'_1 l'_2}^{LS}(r_1, r_2, t).$$

UNCLASSIFIED

TDCC method

- Angular reduction of the two-electron time-dependent Schrödinger equation yields

$$i \frac{\partial}{\partial t} P_{l_1 l_2}^{LS}(r_1, r_2, t) = T_{l_1 l_2} P_{l_1 l_2}^{LS}(r_1, r_2, t) + \sum_{l'_1, l'_2} V_{l_1 l_2, l'_1 l'_2}^L P_{l'_1 l'_2}^{LS}(r_1, r_2, t).$$

- Initial state is a spin-symmetrized product of an incoming electron wavepacket and a valence-shell radial orbital:

$$P_{l_1 l_2}^{LS}(r_1, r_2, t = 0) = \sqrt{\frac{1}{2}} \left[G_{k_0 l_1}(r_1) P_{nl_2}(r_2) + (-1)^S P_{nl_1}(r_1) G_{k_0 l_2}(r_2) \right].$$

UNCLASSIFIED

The challenge of multi-electron atoms

UNCLASSIFIED

The challenge of multi-electron atoms

- Accurate description of inactive multi-electron core and its interaction with active electrons required.

UNCLASSIFIED

The challenge of multi-electron atoms

- Accurate description of inactive multi-electron core and its interaction with active electrons required.
- Higher angular momentum channels must be retained in the wavefunction for a more diffuse atomic system.

UNCLASSIFIED

The challenge of multi-electron atoms

- Accurate description of inactive multi-electron core and its interaction with active electrons required.
- Higher angular momentum channels must be retained in the wavefunction for a more diffuse atomic system.
- Variable radial mesh required to track rapid oscillations of the radial wavefunction near the nucleus.

UNCLASSIFIED

The challenge of multi-electron atoms


- Accurate description of inactive multi-electron core and its interaction with active electrons required.
- Higher angular momentum channels must be retained in the wavefunction for a more diffuse atomic system.
- Variable radial mesh required to track rapid oscillations of the radial wavefunction near the nucleus.
- Retain efficiency and accuracy using an appropriate time propagation scheme.

UNCLASSIFIED

Triple-differential cross sections for electron-impact ionization of Na

UNCLASSIFIED

Manchester experiments




- Detection carried out in coplanar ($\phi_1 = \phi_2 = 0^\circ$) symmetric ($\xi_1 = \xi_2$) or asymmetric geometries.

UNCLASSIFIED

Triple-differential cross sections

coplanar symmetric geometry, equal energy sharing




- Good agreement obtained with experiment in each case; calculations are able to track TDCS features spanning several orders of magnitude.

UNCLASSIFIED

Triple-differential cross sections

coplanar asymmetric geometry, equal energy sharing, $E_{inc} = 11.15$ eV



- Wannier breakup geometry dominates for $\theta_1 = 0^\circ, 150^\circ$.
- Overall preference for $\theta_1 = 30^\circ$.
- Similar trends were observed in previous CCC calculations.


UNCLASSIFIED

Triple-differential cross sections for electron-impact ionization of Mg

UNCLASSIFIED

Triple-differential cross sections

coplanar symmetric geometry, equal energy sharing




- Calculations are able to track TDCS features which range over several orders of magnitude.

UNCLASSIFIED

Triple-differential cross sections

coplanar symmetric geometry, equal energy sharing




- A more complete treatment of valence-shell correlation may be required at 27.65 eV.

UNCLASSIFIED

Triple-differential cross sections

coplanar symmetric geometry, equal energy sharing




- Sensitivity to correlation may be reduced as the 3s electron is ejected more rapidly.

UNCLASSIFIED

Triple-differential cross sections

coplanar asymmetric geometry, equal energy sharing, $E_{inc} = 47.65$ eV




- Good agreement obtained with measurement at $\theta_1 = 30^\circ$.

UNCLASSIFIED

Triple-differential cross sections

coplanar asymmetric geometry, equal energy sharing, $E_{inc} = 47.65$ eV




- Strong preference for emission with $\theta_1 = 30^\circ$, as was the case for Na.

UNCLASSIFIED

Triple-differential cross sections

coplanar asymmetric geometry, equal energy sharing, $E_{inc} = 47.65$ eV



- Additional structure in comparison to the Na TDCS.

UNCLASSIFIED

Summary

- Applied a new version of the TDCC method to calculate angular distributions for ionization of multi-electron targets.
- Good agreement with measurements obtained for a range of incident electron energies for Na and Mg.
- Differing dynamics observed in the angular distributions for He and Mg at a common excess energy.

UNCLASSIFIED

Future plans

- Calculate angular distributions for electron-impact single ionization of Mg where both valence-shell electrons are considered active.

UNCLASSIFIED

Future plans

- Calculate angular distributions for electron-impact single ionization of Mg where both valence-shell electrons are considered active.
- Examine the effect of valence-shell correlation through comparison of two-electron and three-electron calculations.

UNCLASSIFIED

Future plans

- Calculate angular distributions for electron-impact single ionization of Mg where both valence-shell electrons are considered active.
- Examine the effect of valence-shell correlation through comparison of two-electron and three-electron calculations.
- Calculate angular distributions for electron-impact ionization of anisotropic atomic targets and make comparison with recent experiments.

UNCLASSIFIED